

MOTION SICKNESS RELIEF- meclizine hcl tablet

Topco Associates, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topcare 44-403

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

for children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, take the first dose ½ hour to 1 hour before starting activity
- to treat motion sickness, take at first signs of symptoms
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

Questions or comments?

1-888-423-0139

Principal display panel

TopCare
health™

NDC 36800-943-19

COMPARE TO DRAMAMINE® ALL DAY LESS DROWSY ACTIVE INGREDIENT*

LESS DROWSY FORMULA

Motion Sickness Relief

MECLIZINE HCl, 25 mg - ANTIEMETIC

LONG LASTING:

- Up to 24 Hours of Protection
- Prevents Nausea,
Dizziness and Vomiting

**up to
24
HOURS**

•Our Pharmacists Recommend•

8 TABLETS

actual size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**

DISTRIBUTED BY TOPCO ASSOCIATES LLC

ELK GROVE VILLAGE, IL 60007

©TOPCO LNKA0319 QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

*This product is not manufactured or distributed by Medtech Products Inc., owner of
the registered trademark Dramamine® All Day Less Drowsy.

50844 ORG021940319

QUALITY

GUARANTEED



LONG LASTING:

- Up to 24 Hours of Protection
- Prevents Nausea, Dizziness and Vomiting

TopCare
health™

LESS DROWSY FORMULA

Motion Sickness Relief

8 TABLETS

MECLIZINE HCl, 25 mg - ANTIEMETIC

TopCare
health™

NDC 36800-943-19

COMPARE TO DRAMAMINE®
ALL DAY LESS DROWSY
ACTIVE INGREDIENT*

LESS DROWSY FORMULA

Motion Sickness Relief

MECLIZINE HCl, 25 mg - ANTIEMETIC



LONG LASTING:

- Up to 24 Hours of Protection
- Prevents Nausea, Dizziness and Vomiting



8 TABLETS



actual size



E1044

Scan here
for more
information



8-1910-403-19-R
08/02/19-03/19

DISTRIBUTED BY TOPCO ASSOCIATES LLC
 1000 LAKESHORE DRIVE, SUITE 100
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO LKKA0319 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com
 *This product is not manufactured or distributed by Medtronic Products Inc.,
 owner of the registered trademark Dramamine® All Day Less Drowsy.
 150844 086021940319

Questions or comments? 1-888-423-0139


Inactive ingredients corn starch, D&C yellow #10 aluminum lake, lactose anhydrous, magnesium stearate, silicon dioxide

- see end flap for expiration date and lot number
- protect from heat and humidity
- 15°-30°C (59°-86°F)
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Other information
TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- 1/2 hour to 1 hour before starting activity
- to treat motion sickness, take at first signs of symptoms
- adults and children 12 years and over: 1 to 2 tablets once daily, or as directed by a doctor

- Warnings**
- Do not use for children under 12 years of age unless directed by a doctor.
 - Ask a doctor before use if you have:
 - difficulty in urination due to enlargement of the prostate gland
 - glaucoma
 - a breathing problem such as emphysema or chronic bronchitis
 - Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.
 - When using this product:
 - drowsiness may occur
 - avoid alcoholic beverages
 - alcohol, sedatives, and tranquilizers may increase drowsiness
 - use caution when driving a motor vehicle or operating machinery



amish
ate

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

#	Item Code	Package Description	Date	Date
1	NDC:36800-943-19	1 in 1 CARTON	05/30/2019	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final		part336	05/30/2019	

Labeler - Topco Associates, LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(36800-943)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-943)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(36800-943)

Revised: 3/2021

Topco Associates, LLC